



# Simulations and Analytic Models of Relativistic Magnetized Jets

## Citation

Tchekhovskoy, Alexander, Ramesh Narayan, and Jonathan McKinney. 2010. Simulations and Analytic Models of Relativistic Magnetized Jets. In Proceedings of Computational Relativistic Astrophysics: Frontiers of MHD Conference, Princeton, NJ, January 13-16, 2010

## Published Version

<https://www.astro.princeton.edu/~anatoly/PCTS/>

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:35173094>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

# Simulations and Analytic Models of Magnetized Gamma-Ray Burst Jets: Beyond the Progenitor Star

Alexander (Sasha) Tchekhovskoy

with Ramesh Narayan and Jonathan McKinney

Harvard University

Tchekhovskoy et al. (2009, arXiv:0911.2228)

# Gamma-ray bursts

Come in 2 flavors:

Short,  $\lesssim 2$  s



Coalescence of a  
compact object  
binary

Long,  $\gtrsim 2$  s



Death of a  
massive star  
(Woosley 1993)

# Gamma-ray bursts (GRBs)

- **Acceleration**: ultra-relativistic velocity, Lorentz factor  $\gamma \gtrsim 100$  Non-thermal prompt spectrum


- **Collimation**: opening angle  $\theta \lesssim 0.1$
- Relation between **acceleration** and **collimation**:  $\gamma\theta \gtrsim 20$  Jet breaks in afterglow emission

✗ Recent simulations of magnetized (MHD) continuously collimated jets (Komissarov et al. 2009):

$$\gamma\theta \lesssim 1$$


GRB jet quick facts:

1. **Ultra-relativistic**:  $\gamma \gtrsim 100$
2. **Collimated**:  $\theta = 0.04 - 0.2$
3. **Product**  $\gamma\theta \simeq 20 \gg 1$



I will now present the first model of a magnetized GRB jet that correctly reproduces **both collimation** and **acceleration**


# How do magnetic jets work?



# Simulation setup

## Confined Jet

$$\gamma\theta = 2$$



GRB jet quick facts:

1. Ultra-relativistic:  $\gamma \gtrsim 100$
2. Collimated:  $\theta = 0.04 - 0.2$
3. Product  $\gamma\theta \simeq 20 \gg 1$

## Numerical Approach

Time-dependent **ultrarelativistic MHD** equations

Axisymmetry, **perfect** conductivity, and **zero** T

## Problem setup

Perfectly conducting **spinning compact object**

**Collimating wall** of shape  $z \propto R^\alpha$

## Model parameters


Jet wall shape

Spin of compact object

Magnetic field strength

Surface mass loss rate

# Why is $\gamma\theta \lesssim 1$ in confined jets?



- Communication is essential
- Jet boundary B needs to keep announcing its trajectory to the rest of the jet to avoid collisions

GRB jet quick facts:

1. Ultra-relativistic:  $\gamma \gtrsim 100$
2. Collimated:  $\theta = 0.04 - 0.2$
3. Product  $\gamma\theta \simeq 10 \gg 1$

- All signals travel inside Mach cone  $\xi$ :


$$\theta < \xi \approx \frac{1}{\gamma}$$

- Communication across jet  $\rightarrow \theta < \xi$
- Robust conclusion:  $\gamma\theta \lesssim 1$  in confined jets

# Simulation setup


Confined Jet

$$\gamma\theta = 2 \quad \times$$



Deconfined Jet

$$\gamma\theta = 20 \quad \checkmark$$





# Simulation setup

Confined Jet

Deconfined Jet

## Numerically-challenging problem

High magnetization and Lorentz factor ( $\sim 1000$ ):  
very stiff regime

Evolution over 10 orders of magnitude in distance



## Our numerical method uses

Collimating grid that follows field lines at high resolution  
1536x256

Equivalent resolution using non-collimating grid:  
1536x100,000

Evolve only non-stationary region to  
speed up computation


$$\gamma\theta = 20$$



# Confined vs. Deconfined

GRB jet quick facts:


1. Ultra-relativistic:  $\gamma \gtrsim 100$  ✓
2. Collimated:  $\theta = 0.04 - 0.2$  ✓
3. Product  $\gamma\theta \simeq 20 \gg 1$  ✓



**MHD models can produce jet breaks ✓**

# Understand this analytically

After jet loses ambient pressure support, it switches from the **fully confined** solution to the **fully unconfined** solution (AT+ 2009).




$$\begin{aligned}\gamma &= 500 \\ \theta &= 0.04 \\ \gamma\theta &= 20\end{aligned}$$

## GRB jet quick facts:


1. Ultra-relativistic:  $\gamma \gtrsim 100$  ✓
2. Collimated:  $\theta = 0.04 - 0.2$  ✓
3. Product  $\gamma\theta \simeq 20 \gg 1$  ✓

# Understand this analytically (2/3)



# Understand this analytically (3/3)

Centrifugal force slows jet down (approximate)



$$F_m = F_c$$

$$\frac{\epsilon_m}{R} = \frac{\epsilon_m \gamma^2}{R_c}$$

$$\gamma = \left( \frac{R_c}{R} \right)^{1/2}$$

# Required ingredients for GRB jets

Both propagation **inside** and **outside** the star are required for GRB jets:

- 1) Fully **confined** jets are too slow for their opening angles:  $\gamma\theta \lesssim 1$
- 2) Fully **deconfined** jets have too large opening angles

Bottom line: need both

- 1) **confinement** to **collimate** the jet initially and
- 2) **deconfinement** to **accelerate** the jet

# Conclusions

- Numerical & analytical models of magnetized deconfined ultra-relativistic jets, extending over 10 orders of magnitude in distance well into the afterglow region.
- Just outside the star, our jets undergo an abrupt period of acceleration during which  $\gamma$  increases but  $\theta$  is constant
- Deconfinement is necessary to achieve ultrarelativistic  $\gamma$  and  $\gamma\theta \gg 1$  required by jet breaks observations
- Confined jets with subequipartition magnetic fields always have  $\gamma\theta \lesssim 1$
- Future work is the self-consistent simulation of magnetized jet propagation through realistic stellar envelope out to the afterglow region.